

## The general distribution of *Orthoptera* in the main zoogeographical regions of North and Central Asia

Michael G. SERGEEV

Received: 4 May 1993

Accepted for publication: 16 July 1993

SERGEEV M. G. 1993. The general distribution of *Orthoptera* in the main zoogeographical regions of North and Central Asia, Acta zool. cracov. 36(1): 53-76.

**Abstract.** The general distribution of orthopterans is described for the main zoogeographical regions inside the frontiers of the former USSR.

**Key words:** distribution, zoogeography, *Orthoptera*, Palaearctic, regionalization, diversity, endemism.

Michael G. SERGEEV Biological Institute, Siberian Branch of Russian Academy of Sciences, 11, Frunze St., Novosibirsk 630091, Russia; Department of General Biology, Novosibirsk State University 2, Pirogova St., Novosibirsk 630090, Russia.

The scheme of zoogeographical regionalization was proposed for North and Central Asia in earlier work (SERGEEV 1986, 1991, 1992, 1993). I analyzed orthopteran species ranges, compared the position of their boundaries in connection with the ecologic-geographical barriers and tried to classify different boundaries, barriers, and regions.

Now I can propose a division of the Palaearctic Region into the following regions (see also UVAROV 1921; EMELJANOV 1974) (Fig. 1). Most part of these subregions have been described before:

(I) the Eurosiberian Subregion – 2 provinces – about 70 species – a few of them endemics;

(II) the Scythian Subregion – at least 4 provinces – over 200 species – a few of them endemics (mainly montane ones);

(III) the Manchurian Subregion – at least 4 provinces – more than 150 species – many endemic genera and species;

(IV) the Orthrian Subregion – at least 4 provinces – more than 150 species – many endemic genera and species;

(V) the West-Mediterranean Subregion – at least 3 provinces – over 200 species – many endemic genera and species;




Fig. 1. Zoogeographical subregions of the Palearctic Region (see text).

(VI) the East-Mediterranean Subregion – at least 4 provinces – over 200 species – many endemic genera and species;

(VII) the Saharan-Gobian Subregion – at least 22 provinces – more than 600 species – endemic tribes (*Diexini*, *Egnatiini*, *Iranellini*, *Uvaroviini* etc.), many endemic genera and species, especially in the local mountains.

Naturally, these subregions can be divided into provinces, subprovinces, districts etc. The pattern of detailed regionalization has been described previously for North and Middle Asia (mainly the regions inside the frontiers of the former USSR) (SERGEEV 1986, 1988, 1991). A scheme for the well-known European and Mediterranean areas may be worked out in the nearest future. The southern part of the Palearctic Region should be investigated more closely. I think many new taxa will be found in this area, especially in the Orthrian Subregion.

The aim of this paper is to describe the orthopteran species distribution in the main zoogeographical regions of North and Middle Asia and to show the general peculiarities of taxonomic distribution.

The area under study may be divided into the following subregions and provinces (Fig. 2):

(1-2) the Eurosiberian Subregion:

- (1) the Arctic Province,
- (2) the Boreal Province;




Fig. 2. Zoogeographical provinces of North and Central Asia (the Asian part of the former USSR) (see text).

(3-4) the Manchurian Subregion:

- (3) the Amurian Province,
- (4) the Sakhalin-Hokkaido Province;

(5-6) the Scythian Subregion:

- (5) the Russian-Siberian Province,
- (6) the Sarmathian Province;

(7-14) the Saharan-Gobian Subregion:

- (7) the Kazakhstan Province,
- (8) the Mongolian Province,
- (9) the Turanian Province,
- (10) the Gissar-Tien Shan Province,
- (11) the Inner Tien Shan Province,
- (12) the Subpamirian Province,
- (13) the Pamirian Province,
- (14) the Turkmeno-Iranian Province.

The taxonomic composition of each subregion or province is show in the table.



Table I ctd.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<i>Drymadusini</i>														
<i>Lithoxenus grandis</i> (TARBINSKY, 1930)	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>L. nigrofasciatus</i> PRAVDIN, 1979	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>L. heptapotamicus</i> (PYLNOV, 1911)	-	-	-	-	-	-	-	-	-	+	+	-	-	-
<i>L. miramae</i> (VELTISTSHEV, 1940)	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Bergiola balchaschica</i> (STSHELKANOVITZEV, 1907)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>B. montana</i> BEY-BIENKO, 1951	-	-	-	-	-	-	-	-	-	-	-	-	-	+
<i>B. hissarica</i> BEY-BIENKO, 1951	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>B. popovi</i> BEY-BIENKO, 1951	-	-	-	-	-	-	-	-	L	-	-	-	-	-
<i>Eulithoxenus mongolicus</i> (UVAROV, 1928)	-	-	-	-	-	L	-	+	-	-	-	-	-	-
<i>Bienkoxenus beybienkoi</i> (I. STEBAEV, 1964)	-	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Uvarovina daurica</i> (UVAROV, 1928)	-	-	-	-	-	+	-	-	-	-	-	-	-	-
<i>Tadzhikia pavlovskii</i> MISTSHENKO, 1952	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>T. beybienkoi</i> MISTSHENKO, 1952	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ammoxenus pavlovskii</i> BEY-BIENKO, 1951	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>A. desertus</i> BEY-BIENKO, 1951	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Ceraeocercus fuscipennis</i> UVAROV, 1910	-	-	-	-	-	-	+	-	+	+	+	-	-	+
<i>Iranusa khorasana</i> UVAROV, 1942	-	-	-	-	-	-	-	-	+	-	-	-	-	+
<i>Paradrymadusa bucharica</i> PRAVDIN, 1969	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Drymadusella hissarica</i> (MISTSHENKO, 1949)	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Phytodrymadusa longipes</i> (BRUNNER VON WATTENWYL, 1882)	-	-	-	-	-	-	-	-	L <sup>2</sup>	-	-	-	-	-
<i>Ferganusa hemiptera</i> UVAROV, 1939	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Calopterus weneri</i> (ADELUNG, 1910)	-	-	-	-	-	-	-	-	-	-	-	-	-	+
<i>C. pamirica</i> STOLYAROV, 1969	-	-	-	-	-	-	-	-	-	-	-	+	+	-
<i>C. mistshenkoi</i> SERGEEV ET POKIVAJLOV, 1992	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Atlanticus brunneri</i> (PYLNOV, 1914)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Paratlanticus ussuriensis</i> (UVAROV, 1926)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Anatlanticus uvarovi</i> (MIRAM, 1940)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Gampsocleidini</i>														
<i>Gampsocleis glabra</i> (HERBST, 1786)	-	-	-	-	+	+	+	-	+	+	+	-	-	-
<i>G. sedakovii</i> FISCHER DE WALDHEIM, 1846	-	+	+	+	+	+	-	+	-	-	-	-	-	-
<i>G. ussuriensis</i> ADELUNG, 1910	-	-	+	+	-	-	-	-	-	-	-	-	-	-
<i>G. gratiosa</i> BRUNNER VON WATTENWYL, 1888	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>G. shelkownikovae</i> ADELUNG, 1916	-	-	-	-	-	+	-	+	-	-	-	-	-	-
<i>Uvarovites inflatus</i> (UVAROV, 1924)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Decticini</i>														
<i>Decticus verrucivorus</i> (LINNEUS, 1758)	L	+	+	-	+	+	+	+	+	+	+	-	-	+
<i>D. albifrons</i> (FABRICIUS, 1775)	-	-	-	-	-	-	+	-	+	+	-	+	-	+
<i>D. nigrescens</i> TARBINSKY, 1930	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Medecticus assimilis</i> (FIEBER, 1853)	-	-	-	-	-	-	-	-	+	+	-	-	-	+



Table I ctd.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<i>Ctenodecticini</i>														
<i>Miramiola pusilla</i> (MIRAM, 1927)	-	-	-	-	-	+	+	-	-	-	-	-	-	-
<i>Glyphonotinae</i>														
<i>Glyphonotus thoracicus</i> (FISCHER DE WALDHEIM, 1864)	-	-	-	-	-	-	+	-	+	+	-	-	-	+
<i>G. coniciplicus</i> UVAROV, 1914	-	-	-	-	-	-	+	-	+	+	-	-	-	-
<i>G. alajensis</i> MIRAM, 1930	-	-	-	-	-	-	-	-	-	+	-	+	-	-
<i>G. alactaga</i> MIRAM, 1930	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Onconotinae</i>														
<i>Onconotus laxmanni</i> (PALLAS, 1771)	-	-	-	-	+	+	-	-	-	-	-	-	-	-
<i>O. servillei</i> FISCHER DE WALDHEIM, 1846	-	-	-	-	-	+	-	-	-	-	-	-	-	-
<i>Conocephalinae</i>														
<i>Conocephalini</i>														
<i>Conocephalus discolor</i> THUNBERG, 1815	-	-	+	-	+	+	+	-	+	+	-	-	-	-
<i>C. dorsalis</i> (LATREILLE, 1804)	-	-	-	-	+	+	+	-	-	-	-	-	-	-
<i>C. beybienkoi</i> STOROZHENKO, 1981	-	-	+	-	+	+	-	-	-	-	-	-	-	-
<i>C. chinensis</i> (REDTENBACHER, 1891)	-	-	+	+	-	-	-	-	-	-	-	-	-	-
<i>C. japonicus</i> (REDTENBACHER, 1891)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>C. percaudatus</i> BEY-BIENKO, 1955	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>C. buxtoni</i> CHOPARD, 1922	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Copophorini</i>														
<i>Ruspolia nitidula</i> (SCOPOLI, 1786)	-	-	+	-	-	-	+	-	+	-	-	-	-	-
<i>Prophalangopsidae</i>														
<i>Cyphoderrinae</i>														
<i>Paracyphoderris erebeus</i> STOROZHENKO, 1980	-	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>Mimnermidae</i>														
<i>Lezininae</i>														
<i>Lezina mutica</i> (BRUNNER VON WATTENWYL, 1899)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Rhaphidophoridae</i>														
<i>Aemodogryllinae</i>														
<i>Paratachycines boldyrevi</i> (UVAROV, 1926)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>P. ussuriensis</i> STOROZHENKO, 1991	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Diestrarmena unicolor</i> BRUNNER VON WATTENWYL, 1888	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>D. japonica</i> BLATCHLEY, 1920	-	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Gryllidae</i>														
<i>Gryllinae</i>														
<i>Gryllini</i>														
<i>Gryllus bimaculatus</i> DE GEER, 1773	-	-	-	-	-	-	+	-	+	+	-	-	-	+
<i>G. campestris</i> LINNEUS, 1758	-	-	-	-	-	L	L	-	-	-	-	-	-	-
<i>Acheta domestica</i> (LINNEUS, 1758)	-	A <sup>2</sup>	A	-	A	A	A	-	A	-	-	-	-	-
<i>A. turcomana</i> GOROCHOV, 1978	-	-	-	-	-	-	-	-	+	-	-	-	-	L
<i>Melanogryllus desertus</i> (PALLAS, 1771)	-	-	-	-	-	+	+	-	+	+	+	+	+	-
<i>Teleogryllus infernalis</i> (SAUSSURE, 1877)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Tartarogryllus tartarus</i> (SAUSSURE, 1874)	-	-	-	-	-	-	-	-	+	+	-	-	-	+
<i>Modicogryllus frontalis</i> (FIEBER, 1845)	-	-	-	-	-	+	+	+	+	+	-	+	-	-

Table I ctd.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<i>M. pallipalpis</i> (TARBINSKY, 1940)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>M. truncatus</i> (TARBINSKY, 1940)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>M. bordigalensis</i> (LATREILLE, 1804)	-	-	-	-	-	-	+	-	+	+	-	-	-	-
<i>M. chivensis</i> (TARBINSKY, 1930)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>M. bucharicus</i> (BEY-BIENKO, 1933)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Gryllopsis ovtshinnikovi</i> GOROCHOV et MISTSHENKO, 1991	-	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Nigrogryllus sibiricus</i> (CHOPARD, 1925)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Gryllodinus kerkennensis</i> (FINOT, 1893)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>G. odicus</i> (UVAROV, 1911)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>G. abditus</i> GOROCHOV, 1979	-	-	-	-	-	-	-	-	+	-	-	-	-	L
<i>Turanogryllus lateralis</i> (FIEBER, 1853)	-	-	-	-	-	-	-	-	+	+	-	-	-	L
<i>Gryllodes supplicans</i> (WALKER, 1869)	-	-	A	-	-	-	-	-	-	-	-	-	-	-
<i>Loxoblemmus arietulus</i> SAUSSURE, 1877	-	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Velarifictorus micado</i> (SAUSSURE, 1877)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>V. bolivari</i> (UVAROV, 1912)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Cophaphonus riparius</i> MISTSHENKO, 1949	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>C. zimini</i> TARBINSKY, 1932	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>C. tshirkunae</i> MISTSHENKO, 1949	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>C. privatus</i> MISTSHENKO, 1949	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Gryllomorphi</i>														
<i>Gryllomorpha gestroana</i> I. BOLIVAR, 1914	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>G. miramae</i> S. MEDVEV, 1933	-	-	-	-	-	-	-	+	+	-	-	-	-	-
<i>Nemobiinae</i>														
<i>Pteronemobiini</i>														
<i>Pteronemobius nitidus</i> (I. BOLIVAR, 1901)	-	-	+	+	-	-	-	-	-	-	-	-	-	-
<i>P. heydeni</i> (FISCHER, 1853)	-	-	-	-	-	-	-	-	+	+	-	-	-	-
<i>Stenonemobius gracilis</i> (JAKOVLEV, 1871)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Dianemobius fascipes</i> (WALKER, 1869)	-	-	+	+	+	+	-	-	-	-	-	-	-	-
<i>D. furumagiensis</i> (OHMACHI et FURUKAWA, 1929)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>D. csikii</i> (BOLIVAR, 1910)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Polionemobius taprobanensis</i> (WALKER, 1869)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Parapteronemobius sazanami</i> FURUKAWA, 1970	-	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Grylliscini</i>														
<i>Grylliscus gussakovskii</i> TARBINSKY, 1930	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Oecanthinae</i>														
<i>Oecanthus longicaudus</i> MATSUMURA, 1904	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Oe. pellucens</i> (SCOPOLI, 1763)	-	-	-	-	-	+	+	-	+	+	-	+	-	-
<i>Oe. turanicus</i> UVAROV, 1912	-	-	-	-	-	-	-	-	+	+	-	+	-	+
<i>Myrmecophilidae</i>														
<i>Myrmecophilinae</i>														
<i>Myrmecophilini</i>														
<i>Myrmecophilus sapporensis</i> MATSUMURA, 1904	-	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>M. oculatus</i> MIRAM, 1930	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>M. crenatus</i> GOROCHOV, 1986	-	-	-	-	-	-	-	-	+	-	-	-	-	-



Table I ctd.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<i>Bothriophylacini</i>														
<i>Eremogryllodes semenovi</i> (MIRAM, 1930)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>E. vlasovi</i> (MIRAM, 1930)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Gryllotalpidae</i>														
<i>Gryllotalpini</i>														
<i>Gryllotalpa orientalis</i> BURMEISTER, 1838	-	-	+	+	-	-	-	-	+	-	-	-	-	-
<i>G. gryllotalpa</i> (LINNEUS, 1758)	-	-	-	-	-	-	-	-	+?	L2	-	-	-	-
<i>G. stepposa</i> ZHANTIEV, 1991	-	-	-	-	-	-	-	-	+	?	-	-	-	+
<i>G. unispina</i> SAUSSURE, 1874	-	-	-	-	-	+	+	-	+	+	-	-	-	+
<i>Tridactylidae</i>														
<i>Tridactylinae</i>														
<i>Xya variegata</i> LATREILLE, 1809	-	-	-	-	+	+	+	-	+	+	-	-	-	-
<i>X. japonica</i> DE HAAN, 1942	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Tridactylus fasciatus</i> GUERIN-MENEVILLE, 1844	-	-	-	-	-	-	-	-	+	+	-	+	+	-
<i>Dentridactylinae</i>														
<i>Bruntridactylus tartarus</i> (SAUSSURE, 1874)	-	-	-	-	+	+	+	-	+	+	+	+	+	-
<i>Tetrigidae</i>														
<i>Tetriginae</i>														
<i>Tetrix subulata</i> (LINNEUS, 1761)	-	+	+	-	+	+	+	+	+	+	-	-	-	-
<i>T. bolivari</i> SAULCY, 1901	-	-	-	-	-	-	-	-	+	L	-	-	-	-
<i>T. fuliginosa</i> (ZETTERSTEDT, 1828)	+	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>T. tartara</i> (I. BOLIVAR, 1887)	-	-	-	-	+	+	+	+	+	+	+	+	+	+
<i>T. simulans</i> (BEY-BIENKO, 1929)	-	L	+	-	+	+	-	-	-	-	-	-	-	-
<i>T. tenuicornis</i> (SAHLBERG, 1891)	L	+	+	-	+	+	+	+	+	+	-	-	-	-
<i>T. japonica</i> (I. BOLIVAR, 1887)	-	L	+	+	+	+	-	-	-	-	-	-	-	-
<i>T. bipunctata</i> (LINNEUS, 1758)	+	+	+	+	+	+	-	-	-	-	-	-	-	-
<i>Clinotettix ussuriensis</i> BEY-BIENKO, 1933	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Formosatettix robustus</i> STOROZHENKO, 1981	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Paratettix uvarovi</i> SEMENOV, 1915	-	-	-	-	-	-	-	-	+	+	-	+	+	-
<i>P. hachijoensis</i> SHIRAKI, 1906	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Dasyleurotettix depressus</i> (BRISOUT, 1848)	-	-	-	-	-	-	-	-	+	-	-	-	-	+
<i>Ergatettix dorsiferus</i> (WALKER, 1871)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Eumastacidae</i>														
<i>Gomphomastacinae</i>														
<i>Clinomastax ninae</i> (MISTSHENKO, 1937)	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Phytomastax opaca</i> (KRAUSS, 1898)	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ph. artemisiana</i> BEY-BIENKO, 1949	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ph. elegans</i> PRAVDIN, 1969	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ph. hissarica</i> (BEY-BIENKO, 1947)	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ph. robusta</i> (BEY-BIENKO, 1936)	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ph. sijazovi</i> (UVAROV, 1914)	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ph. marikovskij</i> St. TARBINSKIJ, 1955	-	-	-	-	-	-	-	-	-	+	-	-	-	-



Table I ctd.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<i>Pyrgomorphidae</i>														
<i>Pyrgomorphini</i>														
<i>Pyrgomorpha bispinosa</i> WALKER, 1870	-	-	-	-	-	-	+	-	+	+	-	-	-	+
<i>Chrotogonini</i>														
<i>Chrotogonus turanicus</i> KUTHY, 1905	-	-	-	-	-	-	+	-	+	+	+	+	-	-
<i>Acrididae</i>														
<i>Catantopinae</i>														
<i>Uvaroviini</i>														
<i>Uvarovium desertum</i> DIRSH, 1927	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Dericorythini</i>														
<i>Dericorys annulata</i> (FIEBER, 1853)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>D. tibialis</i> (PALLAS, 1773)	-	-	-	-	-	-	-	-	+	L	-	-	-	L
<i>D. albidula</i> AUDINET SERVILLE, 1839	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Egnatiini</i>														
<i>Egnatioides desertus</i> UVAROV, 1926	-	-	-	-	-	-	L	-	+	-	-	-	-	-
<i>Egnatius apicalis</i> STAL, 1876	-	-	-	-	-	-	+	-	+	+	+	-	-	+
<i>Ferganacris mushketovi</i> SERGEEV et BUGROV, 1988	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Diexiini</i>														
<i>Bufonacridella sumakovi</i> ADELUNG, 1910	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Diexis bucharicus</i> MISTSHENKO, 1950	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>D. uvarovi</i> TARBINSKY, 1932	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>D. varentzovi</i> ZUBOVSKY, 1899	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>D. gussakovskiy</i> MIRAM, 1949	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>D. bellus</i> MISTSHENKO, 1950	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>D. chivensis</i> UMN OV, 1931	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>D. ferghanensis</i> UMN OV, 1931	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Iranellini</i>														
<i>Iranella eremiaphila</i> UVAROV, 1922	-	-	-	-	-	-	-	-	L	-	-	-	-	+
<i>I. turcmena</i> BEY-BIENKO, 1948	-	-	-	-	-	-	-	-	L	-	-	-	-	+
<i>Oxyini</i>														
<i>Oxya chinensis</i> (THUNBERG, 1815)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>O. maritima</i> MISTSHENKO, 1951	-	L	+	-	-	-	-	-	-	-	-	-	-	-
<i>O. fuscovittata</i> (MARSCHALL, 1836)	-	-	-	-	-	-	-	-	+	+	-	-	-	-
<i>Tropidopolini</i>														
<i>Tropidopola daurica</i> UVAROV, 1926	-	-	-	-	-	-	?	-	-	-	-	-	-	-
<i>T. turanica</i> UVAROV, 1926	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Conophymatini</i>														
<i>Bienkoa fedtschenkoi</i> (ZUBOVSKY, 1899)	-	-	-	-	-	-	-	-	-	+	-	+	+	-
<i>Conophyma iliense</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>C. leve</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>C. almasyi</i> (KUTHY, 1905)	-	-	-	-	-	-	-	-	-	+	+	-	-	-







Table I ctd.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<i>Acridinae</i>														
<i>Acridini</i>														
<i>Acrida incallida</i> MISTSHENKO, 1951	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>A. cinerea</i> THUNBERG, 1815	-	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>A. oxycephala</i> (PALLAS, 1771)	-	-	-	-	-	-	+	-	+	+	-	-	+	+
<i>Truxalini</i>														
<i>Truxalis eximia</i> (EICHWALD, 1830)	-	-	-	-	-	-	-	-	+	-	-	-	-	+
<i>Ochrilidini</i>														
<i>Gonista sagitta</i> (UVAROV, 1912)	-	-	-	-	-	-	-	-	+	L	-	-	-	L
<i>Ochrilidia mistshenkoi</i> (BEY-BIENKO, 1936)	-	-	-	-	-	-	-	-	+	+	-	-	-	-
<i>O. turanica</i> (BEY-BIENKO, 1936)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>O. hebetata</i> (UVAROV, 1927)	-	-	-	-	-	-	-	-	+	+	-	-	-	-
<i>Phlaeobini</i>														
<i>Duroniella brachyptera</i> UMNOV, 1931	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>D. kalmyka</i> (ADELUNG, 1906)	-	-	-	-	-	-	+	-	+	L	-	-	-	L
<i>D. gracilis</i> UVAROV, 1926	-	-	-	-	-	-	+	-	+	+	-	-	-	+
<i>D. turcomana</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>D. sogdiana</i> MISTSHENKO, 1949	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Chrysochraontini</i>														
<i>Chrysochraon dispar</i> (GERMAR, 1831)	-	+	+	+	+	+	-	-	-	+	-	-	-	-
<i>Ch. amurensis</i> MISTSHENKO, 1989	-	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>Euthystira brachyptera</i> (OCSKAY, 1826)	-	+	+	+	+	+	+	-	-	+	+	-	-	-
<i>Eu. japonica</i> (I. BOLIVAR, 1898)	-	-	+	+	+	+	-	+	-	-	-	-	-	-
<i>Podismopsis altaica</i> ZUBOVSKY, 1900	-	-	-	-	+	+	L	-	-	-	-	-	-	-
<i>P. gynaemorpha</i> IKONNIKOV, 1911	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>P. genicularibus</i> (SHIRAKI, 1910)	-	+	+	+	-	-	-	-	-	-	-	-	-	-
<i>P. poppiusi</i> (MIRAM, 1907)	L	+	-	-	+	L	-	-	-	-	-	-	-	-
<i>P. jacuta</i> MIRAM, 1928	-	+	-	-	L	-	-	-	-	-	-	-	-	-
<i>P. ussuriensis</i> IKONNIKOV, 1911	-	-	+	-	+	-	-	-	-	-	-	-	-	-
<i>P. gelida</i> MIRAM, 1931	L	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>P. insularis</i> MISTSHENKO, 1951	-	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>P. silvestris</i> STOROZHENKO, 1981	-	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>P. konakovi</i> BEY-BIENKO, 1948	-	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Hypernephiini</i>														
<i>Eclipophleps beybienkoi</i> MALJKOVSKI, 1972	-	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>E. kazacha</i> MALJKOVSKI, 1959	-	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>E. glacialis</i> BEY-BIENKO, 1933	-	-	-	-	-	+	-	-	-	-	-	-	-	-
<i>Saxetophilus petulans</i> UMNOV, 1930	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>S. mistshenkoi</i> NAUMOVICH, 1984	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Arcypterini</i>														
<i>Arcyptera fusca</i> (PALLAS, 1773)	-	+	-	-	+	+	L	L	-	+	-	-	-	-
<i>A. orientale</i> STOROZHENKO, 1990	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Pararcyptera microptera</i> (FISCHER DE WALDHEIM, 1833)	-	+	+	-	+	+	+	+	L	+	+	-	-	+

Table I ctd.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<i>Ramburiella turcomana</i> (FISCHER DE WALDHEIM, 1833)	-	-	-	-	-	-	-	-	+	+	+	+	-	+
<i>R. foveolata</i> TARBINSKY, 1931	-	-	-	-	-	-	-	-	+	+	-	-	+	+
<i>R. bolivari</i> (KUTHY, 1907)	-	-	-	-	-	-	-	-	+	+	+	+	-	+
<i>Dociostaurini</i>														
<i>Dociostaurus maroccanus</i> (THUNBERG, 1815)	-	-	-	-	-	-	-	-	+	+	-	-	-	+
<i>D. brevicollis</i> (EVERSMANN, 1848)	-	-	-	-	+	+	+	-	+	+	+	-	+	-
<i>D. tartarus</i> (STSHELKANOVITZEV, 1909)	-	-	-	-	-	-	+	-	+	+	+	+	+	+
<i>D. plotnikovi</i> UVAROV, 1921	-	-	-	-	-	-	-	-	+	L	-	-	-	L
<i>D. kraussi</i> (INGENENITZKY, 1897)	-	-	-	-	-	+	+	-	+	+	+	+	-	+
<i>Notostaurus albicornis</i> (EVERSMANN, 1848)	-	-	-	-	-	+	+	-	+	+	+	+	+	+
<i>N. popovi</i> MIRAM, 1935	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Kazakia tarbinskyi</i> BEY-BIENKO, 1933	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Mizonocara uvarovi</i> BEY-BIENKO, 1933	-	-	-	-	-	-	-	-	L	+	-	-	-	-
<i>M. kusnezovae</i> UMNNOV, 1947	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>M. notata</i> MISTSHENKO, 1947	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>M. robusta</i> MISTSHENKO, 1947	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>M. inornata</i> MISTSHENKO, 1947	-	-	-	-	-	-	-	-	-	-	-	-	-	+
<i>M. deserti</i> UVAROV, 1912	-	-	-	-	-	-	-	-	-	-	-	-	-	+
<i>M. saksinae</i> MISTSHENKO, 1987	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Eremippus persicus</i> UVAROV, 1929	-	-	-	-	-	-	-	-	+	L	-	-	-	-
<i>E. comatus</i> MISTSHENKO, 1951	-	-	-	-	-	-	+	-	+	-	-	-	-	-
<i>E. carinatus</i> MISTSHENKO, 1951	-	-	-	-	-	-	L	-	+	-	-	-	-	-
<i>E. mistshenkoi</i> I. STEBAEV, 1965	-	-	-	-	-	-	+	+	-	-	?	-	-	-
<i>E. simplex</i> (EVERSMANN, 1859)	-	-	-	-	-	L	+	+	+	+	-	-	-	+
<i>E. nudus</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	-	-	-	+	-
<i>E. miramae</i> TARBINSKY, 1940	-	-	-	-	-	L	+	-	+	+	-	-	-	L
<i>E. foveolatus</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	+	+	-	-	-	-
<i>E. beybienkoi</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>E. veltistishevi</i> MIRAM, 1935	-	-	-	-	-	-	-	-	+	-	+	-	-	-
<i>E. onerosus</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	-	-	-	-	+
<i>E. betpakdalensis</i> SKOPIN, 1951	-	-	-	-	-	-	+	-	+	-	-	-	-	-
<i>E. selevini</i> SKOPIN, 1951	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>E. flavus</i> MISTSHENKO, 1951	-	-	-	-	-	-	?	-	-	-	-	-	-	-
<i>E. pusillus</i> BEY-BIENKO, 1948	-	-	-	-	-	-	+	-	+	-	-	-	-	-
<i>E. pilosus</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>E. luppovae</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	-	-	-	+	-
<i>E. hemipterus</i> MALJKOVSKI, 1968	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>E. nanus</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Gomphocerini</i>														
<i>Stenobothrus lineatus</i> (PANZER, 1796)	-	+	+	-	+	+	+	-	-	+	-	-	-	-
<i>S. olgaephilus</i> STOROZHENKO, 1985	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>S. fischeri</i> (EVERSMANN, 1848)	-	-	-	-	+	+	+	-	+	+	+	-	-	-
<i>S. miramae</i> DIRSH, 1931	-	-	-	-	-	+	+	-	-	-	-	-	-	-



Table I ctd.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<i>S. nigromaculatus</i> (HERICH-SCHAFFER, 1840)	-	L	-	-	+	+	+	-	-	+	+	-	-	-
<i>S. kirgisorum</i> IKONNIKOV, 1911	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>S. cobresianus</i> BEY-BIENKO, 1949	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>S. carbonarius</i> (EVERSMANN, 1848)	-	-	-	-	-	+	+	+	+	+	-	-	-	-
<i>S. newskii</i> ZUBOVSKY, 1899	-	-	-	-	+	+	-	-	-	-	-	-	-	-
<i>S. eurasius</i> ZUBOVSKY, 1898	-	L	-	-	+	+	+	+	+	+	+	-	-	-
<i>S. tadzhicus</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Omocestus viridulus</i> (LINNEUS, 1758)	-	+	+	+	+	+	+	+	-	+	+	-	-	-
<i>O. heymonsi</i> (RAMME, 1926)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>O. ventralis</i> (ZETTERSTEDT, 1821)	-	?	?	-	?	?	?	-	?	-	-	-	-	-
<i>O. haemorrhoidalis</i> (CHARPENTIER, 1825)	-	+	+	-	+	+	+	+	+	+	+	+	-	-
<i>O. petraeus</i> (BRISOUT-BARNEVILLE, 1855)	-	-	-	-	+	+	+	-	+	L	-	-	-	-
<i>Myrmeleotettix palpalis</i> (ZUBOVSKY, 1899)	-	-	L	-	+	+	L	+	-	-	-	-	-	-
<i>M. antennatus</i> (FIEBER, 1853)	-	-	-	-	-	+	+	-	L	-	-	-	-	-
<i>M. maculatus</i> (THUNBERG, 1815)	-	-	-	-	+	+	-	-	-	-	-	-	-	-
<i>M. pallidus</i> (BRUNNER VON WATTENWYL, 1882)	-	-	-	-	+	+	+	-	+	-	-	-	-	-
<i>Gomphocerus rufus</i> (LINNEUS, 1758)	-	+	L	-	+	+	-	-	-	-	-	-	-	-
<i>Aeropus kudia</i> CAUDELL, 1927	-	+	+	-	-	-	-	-	-	-	-	-	-	-
<i>Ae. sibiricus</i> (LINNEUS, 1767)	+	+	L	-	+	+	+	-	-	+	+	+	+	-
<i>Aeropedellus variegatus</i> (FISCHER DE WALDHEIM, 1846)	+	+	-	-	+	+	-	-	-	-	-	-	-	-
<i>Ae. reuteri</i> (MIRAM, 1907)	-	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Ae. baliolus</i> MISTSHENKO, 1951	-	-	-	-	-	+	-	-	-	-	-	-	-	-
<i>Ae. volgensis</i> (PREDTETSHENSKY, 1928)	-	-	-	-	-	-	+	-	+	-	-	-	-	-
<i>Dasyhippus barbipes</i> (FISCHER DE WALDHEIM, 1846)	-	-	-	-	L	+	-	+	-	-	-	-	-	-
<i>Mesasippus divergens</i> (BEY-BIENKO, 1930)	-	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>M. geophilus</i> (BEY-BIENKO, 1948)	-	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>M. tarbagataicus</i> SERGEEV et BUGROV, 1988	-	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>M. fuscovittatus</i> (TARBINSKY, 1927)	-	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>M. scitus</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>M. ammophilus</i> BEY-BIENKO, 1948	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>M. barsukiensis</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>M. arenosus</i> (BEY-BIENKO, 1930)	-	-	-	-	-	+	+	-	-	-	-	-	-	-
<i>M. kozhevnikovi</i> (TARBINSKY, 1925)	-	-	-	-	-	-	+	-	+	-	-	-	-	-
<i>M. nudus</i> (UMNOV, 1931)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Pezohippus callosus</i> (UVAROV, 1926)	-	-	-	-	-	-	-	-	+	L	-	-	-	-
<i>Stauroderus scalaris</i> (FISCHER DE WALDHEIM, 1846)	-	+	-	-	+	+	+	-	+	+	+	+	-	-
<i>Megaulacobothrus aethalinus</i> (ZUBOVSKY, 1899)	-	L	+	-	+	-	-	-	-	-	-	-	-	-
<i>Glyptobothrus brunneus</i> (THUNBERG, 1815)	-	?	?	-	?	?	?	-	?	?	-	-	-	?
<i>G. miramae</i> RAMME, 1939	-	-	-	-	-	L	-	-	-	-	-	-	-	-
<i>G. biguttulus</i> (LINNEUS, 1758), s. l.	L	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>G. mollis</i> (CHARPENTIER, 1825)	-	-	-	-	+	+	+	-	-	+	-	-	-	+
<i>G. dubius</i> (ZUBOVSKY, 1898)	-	-	-	-	+	+	+	+	-	-	-	-	-	-

Table I ctd.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<i>G. hemipterus</i> UVAROV, 1926	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Chorthippus vagans</i> (EVERSMANN, 1848)	-	-	-	-	-	+	+	-	+	-	-	-	-	-
<i>Ch. uvarovi</i> (BEY-BIENKO, 1929)	-	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Ch. pilipes</i> BEY-BIENKO, 1933	-	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Ch. apriciarius</i> (LINNEUS, 1758)	-	+	-	-	+	+	+	-	-	+	+	+	+	+
<i>Ch. intermedius</i> (BEY-BIENKO, 1926)	-	+	+	-	+	+	-	+	-	-	-	-	-	-
<i>Ch. hammarstroemi</i> (MIRAM, 1907)	-	+	+	-	+	+	-	+	-	-	-	-	-	-
<i>Ch. macrocerus</i> (FISCHER DE WALDHEIM, 1846)	-	-	-	-	-	+	+	-	+	-	-	-	-	+
<i>Ch. saxatilis</i> BEY-BIENKO, 1948	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. ketmenicus</i> BEY-BIENKO, 1949	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. ingenitzkyi</i> (ZUBOVSKY, 1898)	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. kusnetzovi</i> BEY-BIENKO, 1949	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. cavilosus</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. similis</i> UMNNOV, 1930	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. tianschanicus</i> UMNNOV, 1930	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. oreophilus</i> BEY-BIENKO, 1948	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. monilicornis</i> UMNNOV, 1931	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. antennalis</i> UMNNOV, 1931	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. vicinus</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	+	+	+	-	-
<i>Ch. kirgizicus</i> MISTSHENKO, 1979	-	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Ch. shantariensis</i> MISTSHENKO, 1951	-	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>Ch. robustus</i> MISTSHENKO, 1979	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. ferghanensis</i> UMNNOV, 1931	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. karateghinicus</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Ch. darvazicus</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Ch. maracandicus</i> MISTSHENKO, 1979	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. transalajicus</i> MISTSHENKO, 1979	-	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Ch. zaitzevi</i> MISTSHENKO, 1979	-	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Ch. badius</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. plotnikovi</i> UMNNOV, 1931	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. tadzhicus</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Ch. pavlovskii</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. jachontovi</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. jacobsoni</i> (IKONNIKOV, 1911)	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. bucharicus</i> BEY-BIENKO, 1948	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. curtus</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. pascuus</i> UMNNOV, 1931	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Ch. fallax</i> (ZUBOVSKY, 1900)	-	+	+	+	+	+	-	+	-	-	-	-	-	-
<i>Ch. parallelus</i> (ZETTERSTEDT, 1821)	-	+	-	-	+	+	+	-	+	+	-	-	-	-
<i>Ch. turanicus</i> TARBINSKY, 1927	-	-	-	-	-	-	-	-	+	+	-	-	-	-
<i>Ch. montanus</i> (CHARPENTIER, 1825)	L	+	+	+	+	+	-	+	-	-	-	-	-	-
<i>Ch. dorsatus</i> (ZETTERSTEDT, 1821)	-	-	+	-	+	+	L?	-	-	-	-	-	-	-

Table I ctd.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<i>Ch. dichrous</i> (EVERSMANN, 1859)	-	-	-	-	-	+	+	+	+	+	-	+	-	-
<i>Ch. karatavicus</i> BEY-BIENKO, 1936	-	-	-	-	-	-	-	-	L	+	-	-	-	-
<i>Ch. luminosus</i> MISTSHENKO, 1951	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Ch. albomarginatus</i> (DE GEER, 1773)	-	+	+	-	+	+	+	+	+	+	+	-	-	-
<i>Ch. angulatus</i> TARBINSKY, 1927	-	-	-	-	-	-	-	-	+	+	+	-	-	-
<i>Ch. schmidti</i> (IKONNIKOV, 1913)	-	-	+	-	+	+	-	-	-	-	-	-	-	-
<i>Euchorthippus unicolor</i> (IKONNIKOV, 1913)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Eu. pulvinatus</i> (FISCHER DE WALDHEIM, 1846)	-	-	-	-	-	+	+	-	+	+	+	+	-	-
<i>Eu. transcaucasicus</i> TARBINSKY, 1930	-	-	-	-	-	-	-	-	-	-	-	-	-	+
<i>Mecostethini</i>														
<i>Stethophyma grossum</i> (LINNEUS, 1758)	L	+	+	-	+	+	-	+	-	-	-	-	-	-
<i>S. tsherskii</i> IKONNIKOV, 1911	-	+	+	+	+	+	-	-	-	-	-	-	-	-
<i>Mecostethus alliaceus</i> (GERMAR, 1817)	-	-	+	-	+	+	+	-	+	+	-	-	-	-
<i>Epacromiini</i>														
<i>Epacromius pulverulentus</i> (FISCHER DE WALDHEIM, 1846)	-	-	+	-	+	+	+	+	L	+	-	-	-	-
<i>E. tergestinus</i> (CHARPENTIER, 1815)	-	-	-	-	+	+	+	+	+	+	+	+	+	-
<i>Aiolopus thalassinus</i> (FABRICIUS, 1781)	-	-	?	-	-	+	+	-	+	+	-	-	+	+
<i>A. oxianus</i> UVAROV, 1926	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>A. simulatrix</i> (WALKER, 1870)	-	-	-	-	-	-	-	-	+	L	-	-	-	+
<i>A. strepens</i> (LATREILLE, 1804)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Hilethera turanica</i> UVAROV, 1925	-	-	-	-	-	-	-	-	-	+	+	-	-	L
<i>Locustini</i>														
<i>Locusta migratoria</i> LINNEUS, 1758	-	-	+	+	M	+	+	-	+	+	-	-	+	L
<i>Oedaleus infernalis</i> SAUSSURE, 1884	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Oe. decorus</i> (GERMAR, 1817)	-	-	-	-	+	+	+	+	+	+	+	+	+	+
<i>Oe. senegalensis</i> (KRAUSS, 1877)	-	-	-	-	-	-	-	-	+	+	-	-	-	-
<i>Psophus stridulus</i> (LINNEUS, 1758)	-	+	+	-	+	+	-	-	-	-	-	-	-	-
<i>Pyrgodera armata</i> FISCHER DE WALDHEIM, 1846	-	-	-	-	-	-	+	-	+	+	-	-	-	+
<i>Brunnerella mirabilis</i> SAUSSURE, 1888	-	-	-	-	-	-	-	-	-	-	-	-	-	+
<i>Ptetica cristulata</i> SAUSSURE, 1884	-	-	-	-	-	-	-	-	+	+	-	-	-	-
<i>Oedipodini</i>														
<i>Celes variabilis</i> (PALLAS, 1774)	-	L	-	-	+	+	+	-	+	+	+	-	-	-
<i>C. skalozubovi</i> ADELUNG, 1906	-	-	+	-	+	+	-	+	-	-	-	-	-	-
<i>Mioscirtus wagneri</i> (EVERSMANN, 1846)	-	-	-	-	-	-	+	-	+	+	-	-	-	L
<i>Oedipoda juxartensis</i> UVAROV, 1912	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Oe. himalayana</i> UVAROV, 1925	-	-	-	-	-	-	-	-	-	-	-	-	+	-
<i>Oe. fedtschenkoi</i> SAUSSURE, 1884	-	-	-	-	-	-	-	-	-	+	-	+	+	-
<i>Oe. caerulescens</i> (Linneus, 1758)	-	-	-	-	A	+	+	-	+	+	+	-	-	-
<i>Oe. miniata</i> (PALLAS, 1771)	-	-	-	-	-	+	+	-	+	+	+	-	-	+
<i>Trilophidiini</i>														
<i>Trilophidia annulata</i> (THUNBERG, 1815)	-	-	+	-	-	-	-	-	-	-	-	-	-	-
<i>Acrotlyini</i>														
<i>Acrotylus insubricus</i> (SCOPOLI, 1786)	-	-	-	-	-	-	-	+	-	+	+	-	-	-

Table I ctd.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<i>Bryodemini</i>														
<i>Bryodema holdereri</i> KRAUSS, 1901	-	L	-	-	+	+	-	+	-	-	-	-	-	-
<i>B. tuberculatum</i> (FABRICIUS, 1775)	-	+	+	-	+	+	L	-	-	-	-	-	-	-
<i>B. zaisanicum</i> BEY-BIENKO, 1930	-	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>B. orientale</i> BEY-BIENKO, 1930	-	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>B. semenovi</i> IKONNIKOV, 1911	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>B. gebleri</i> (FISCHER DE WALDHEIM, 1836)	-	-	-	-	L	+	+	+	-	+	+	-	-	-
<i>B. heptapotamicum</i> BEY-BIENKO, 1930	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>B. luctuosum</i> (STOLL, 1813)	-	-	-	-	+	+	-	-	-	-	-	-	-	-
<i>Angaracris barabensis</i> (PALLAS, 1773)	-	-	L	-	+	+	+	+	-	-	-	-	-	-
<i>A. rhodopa</i> (FISCHER DE WALDHEIM, 1836)	-	-	-	-	-	+	-	+	-	-	-	-	-	-
<i>Compsorhhipis davidiana</i> (SAUSSURE, 1888)	-	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Sphingonotini</i>														
<i>Pseudococeles persa</i> (SAUSSURE, 1884)	-	-	-	-	-	-	-	-	-	-	-	-	-	+
<i>Cophotylus decorus</i> BEY-BIENKO, 1951	-	-	-	-	-	-	-	-	-	-	-	-	+	-
<i>Sphingonotus halophilus</i> BEY-BIENKO, 1929	-	-	-	-	-	-	+	-	+	-	-	-	-	-
<i>S. kirgizicus</i> MISTSHENKO, 1936	-	-	-	-	-	-	-	-	-	+	+	+	-	-
<i>S. pamiricus</i> RAMME, 1930	-	-	-	-	-	-	-	-	-	+	-	-	+	-
<i>S. zebra</i> MISTSHENKO, 1936	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>S. miramae</i> MISTSHENKO, 1936	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>S. turcmenus</i> BEY-BIENKO, 1951	-	-	-	-	-	-	-	-	-	-	-	-	-	+
<i>S. maculatus</i> UVAROV, 1925	-	-	-	-	-	-	+	-	+	+	+	+	+	+
<i>S. halocnemi</i> UVAROV, 1925	-	-	-	-	-	-	-	-	+	L	-	-	-	-
<i>S. rubescens</i> (WALKER, 1870)	-	-	-	-	-	-	+	-	+	+	+	+	+	+
<i>S. pilosus</i> SAUSSURE, 1884	-	-	-	-	-	-	-	-	+	+	-	-	-	+
<i>S. elegans</i> MISTSHENKO, 1936	-	-	-	-	-	-	-	-	+	+	+	+	+	+
<i>S. lucidus</i> MISTSHENKO, 1936	-	-	-	-	-	-	-	-	-	-	-	-	+	-
<i>S. beybienkoi</i> MISTSHENKO, 1936	-	-	-	-	-	-	+	+	+	-	-	-	-	-
<i>S. caeruleans</i> (LINNEUS, 1767)	-	-	-	-	-	-	+	-	+	L	-	-	-	-
<i>S. coeruleipes</i> (UVAROV, 1922)	-	-	-	-	-	L	+	-	+	+	-	-	-	-
<i>S. eurasius</i> MISTSHENKO, 1936	-	-	-	-	-	-	+	-	+	L	-	-	-	+
<i>S. nebulosus</i> (FISCHER DE WALDHEIM, 1846)	-	-	-	-	-	-	+	-	+	+	+	+	+	+
<i>S. mongolicus</i> SAUSSURE, 1888	-	-	+	-	+	+	-	-	-	-	-	-	-	-
<i>S. octofasciatus</i> (AUDINET SERVILLE, 1839)	-	-	-	-	-	-	-	-	+	+	-	-	+	+
<i>S. obscuratus</i> (WALKER, 1925)	-	-	-	-	-	-	-	-	+	L	-	-	-	L
<i>S. satrapes</i> SAUSSURE, 1884	-	-	-	-	-	-	-	-	+	L	-	-	-	L
<i>S. salinus</i> (PALLAS, 1773)	-	-	-	-	-	-	+	+	+	-	-	-	-	-
<i>Pseudosphingonotus savignyi</i> (SAUSSURE, 1884)	-	-	-	-	-	-	-	-	+	L	-	-	-	+
<i>Sphingoderus carinatus</i> (SAUSSURE, 1888)	-	-	-	-	-	-	L	-	+	L	-	-	-	+
<i>Helioscirtus moseri</i> SAUSSURE, 1884	-	-	-	-	-	-	-	-	+	-	-	-	-	L
<i>Hyalorrhhipis clausi</i> (KITARY, 1849)	-	-	-	-	-	-	+	-	+	-	-	-	-	-
<i>H. turcmena</i> UVAROV, 1926	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Leptopternis gracilis</i> (EVERSMANN, 1848)	-	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>L. iliensis</i> UVAROV, 1925	-	-	-	-	-	-	-	-	+	-	-	-	-	-

The katydids of the subfamily *Deracanthinae* (*Bradyporidae*) are mainly Mongolian and Chinese species. Their numerous endemic genera and species are distributed throughout the north-eastern provinces of the Saharan-Gobian Subregion. As a rule, the *Deracanthini* occur in the northern parts of this area, and the *Zichyini* are connected with its southern parts. The *Phaneropterinae* are chiefly associated with subtropical and tropical regions. Some of them are found in the study area. Most short-winged *Odonturini* (especially *Isophya* and *Poecilimon*) may be characterized as endemic or subendemic. *Saga pedo* is the sole species of the *Saginae* in Palaearctic Asia.

The *Tettigoniinae* katydids are very various. Four species of the *Tettigoniini* proper are distributed mainly throughout the southern part of the forest zone, the forest-steppe and steppe zones. The *Drymadusini* include many species with small ranges. Some of them are endemic in different montane ridges. At the same time, many species are associated with arid areas or with nemoral regions of the Far East. The tribes *Gampsocleidini* and *Decticini* are widely distributed all over the Palearctic. They include some endemic species of the Manchurian Subregion. The distribution of *Platycleidini* is similar to that of the *Drymadusini* but the species of the former are as a rule, comparatively widely distributed. It is important that this tribe includes some forms which are connected with the forest-steppe and steppe zones (*Montana striata*, *M. tomini* etc.) or the forest one (*Roeseliana roeseli*). There are many montane endemics too. The only species of the *Ctenodecticini* occurs within the boundaries of steppes and semi-deserts.

The subfamily *Glyphonotinae* is endemic in the central part of the Saharan-Gobian Subregion. It consists of plain-montane and montane forms. The very specific subfamily *Onconotinae* spreads throughout the Sarmathian Province. Only *Hyphinomos* is supposed to be a Tibetan member of this group. The *Conocephalinae* katydids are chiefly connected with tropical regions. But some species of two tribes – *Conocephalini* and *Copiphorini* – are distributed in the southern part of the Palaearctic. As a rule, they do not penetrate into the Eurosiberian Subregion.

Three families of camel-cricket – *Prophalangopsidae*, *Mimnermidae*, and *Rhaphidophoridae* – include a few Palaearctic species. The sole species of *Paracyphoderris* occurs in the forest part of the Lower Amur Basin. *Lezina mutica* from the *Mimnermidae* does not penetrate from the south into the northern deserts. Some species of the *Rhaphidophoridae* are limited by the Manchurian Subregion.

In the area studied crickets from the family *Gryllidae* are mainly associated with the Manchurian (especially *Nemobiini*) and Saharan-Gobian (*Gryllini*) subregions. But they usually inhabit river and lake valleys. A few forms penetrate into the Scythian Subregion. The *Myrmecophilidae* and *Gryllotalpidae* comprise also some species of the Manchurian and Saharan-Gobian subregion. Their distribution is not well known.

Most of the Palaearctic *Tridactylidae* and *Tetrigidae* are connected with the forest and forest-steppe zones. The former are distributed in the southern part of the study area, and the latter penetrate into the tundra zone. But some species resemble the crickets in being widely distributed in river valleys of an arid area.

The *Gomphomastacinae* (*Eumastacidae*) are an endemic group in the mountains of Tien Shan, Pamiro-Allay, Nanshan, Karakoram, the Hindu-Kush and Himalayas. Most species live at closely localized sites, lying in a definite part of each ridge.

The grasshoppers proper (*Pamphagidae*, *Pyrgomorphidae*, and *Acrididae*) are the main group of the Palaearctic *Orthoptera*. But two species of the *Pyrgomorphidae* are distributed in Middle Asia. They do not penetrate through the northern boundaries of the semi-deserts.

The *Pamphagidae* include forms common to the arid plains and mountains of Middle Asia. There are many endemic genera and species, which may be connected with different types of deserts and semi-deserts (stone, clay, sand). Some species are distributed throughout the Sarmathian Province. The only endemic form in the Amur Basin and Manchuria is *Haplotropis brunneriana*. It is important that its relatives inhabit the Mediterranean area.

The family *Acrididae* comprises at least two subfamilies – *Catantopinae* and *Acridinae*. Some *Catantopinae* tribes are endemics in the Saharan-Gobian Subregion (*Uvaroviini*, *Dericorythini*, *Egnatiini*, *Diexini*, *Iranellini*). The *Conophymatini* grasshoppers live in the mountains of Middle Asia, including the mountains of Iran, Afghanistan, and the West Himalayas. They include many wingless endemics, which are chiefly associated with the outer ridges of these mountains. The Palaearctic *Podismini* (= *Melanoplini*) partly resemble their North American relatives. As a rule, they are short-winged herbicolous grasshoppers, but such forms may be more or less widely distributed. At the same time, many *Primnoa* species are endemics in the Manchurian Subregion. The sole species of the *Cyrthacanthacridini* inhabits Middle Asia. But the Desert Locust is able to penetrate through the frontiers of the former USSR. On the contrary, the *Calliptamini* are characteristic forms of the steppe, semi-desert and desert landscapes of the study area. They often dominate in the local orthopteran communities. The *Eyprepocnemidini* are limited to the Saharan-Gobian and Manchurian subregions.

A few tribes of the *Acridinae* (*Acridini*, *Truxalini*, *Ochrlidini*, *Phlaeobini*, *Acrotylini*) are chiefly connected with the Saharan-Gobian Subregion and occur, as a rule, in local azonal landscapes. The *Chrysochraontini* are limited by the Holarctic boundaries. They include short-winged species. Some of them are widely distributed. But most Palaearctic species are endemic in the Far East. The distribution of the *Hypernephiini* is very interesting. These short-winged forms inhabit mainly the arid mountains of East Kazakhstan and Mongolia. Some genera are limited to the mountains of Tien Shan, Zagros, the East Himalayas and Hengduanshan. The *Arcypterini* and *Gomphocerini* are distributed all over the Holarctic area. They include both widely distributed forms of plains and highly localized montane species. The *Dociostaurini* are connected with the arid areas of the Scythian and Saharan-Gobian subregions. Some of them are provincial endemics.

Most species of the mainly tropical and subtropical *Epacromiini*, *Locustini*, *Oedipodini*, and *Trilophidiini* are limited to the southern part of study area. The mecostethine grasshoppers are associated with the forest and steppe zones of the Holarctic. The *Bryodemini* and *Sphingonotini* consist of terricolous grasshoppers, which occupy mainly the Saharan-Gobian Subregion. But the former is chiefly limited to Central Asia proper

(it should be emphasized that the North American genera *Trimerotropis* and *Circotettix* do not belong to this tribe). The latter is widely distributed throughout the arid regions of the world.

On the whole, it is evident that the pattern of distribution results from the interaction of both the present natural (geographical, ecological, climatic etc. ) conditions and the evolutionary history of the spread of species. Generally speaking, the *Orthoptera* distribution in North and Middle Asia reflects the southern thermophilic character of this group as well as the connection of most its species with grassland and arid ecosystems.

The majority of higher taxa consist of tropical and subtropical groups. Only a few subfamilies (*Deracanthinae*, *Glyphonotinae*, *Onconotinae*, *Thrinchinae*, *Pamphaginae*) and tribes (*Odonturini*, *Drymadusini*, *Gampsocleidini*, *Uvaroviini*, *Egnatiini*, *Conophymatini*, *Chrysochraontini* etc.) are mainly connected with the Palaearctic Region, but almost all the taxa have their centres of diversity and endemism in the southern part of that area. Some of them are associated with local mountains.

An analysis of the distribution of the centres of diversity and generic endemism carried out for each tribe allows us to understand the pattern, which reflects the general situation described above on the Earth's surface. Most centres of tribal diversity and generic endemism are connected with 4 areas of North and Middle Asia:

I – the deserts of Turan (*Drymadusini*, *Diexini*, *Dociostaurini* etc.; endemic – *Grylliscus*);

II – the arid part of Mongolia and China (*Deracanthini*, *Drymadusini*, *Gomphocerini*, *Bryodemini* etc.; endemics – *Bienkoxenus*, *Eulithoxenus*, *Pseudotmethis*, *Sinotmethis* etc.);

III – the nemoral regions near the boundary of North and East Asia (*Meconematinae*, *Drymadusini*, *Gampsocleidini*, *Podismini* etc.; endemics – *Uvarovites*, *Nigrogryllus*, *Clinotettix* etc.);

IV – the mountains of Tien Shan, Pamiro-Allay, and the Hindu-Kush in Afghanistan (*Gomphomastacinae*, *Drymadusini*, *Platycleidini*, *Egnatiini*, *Conophymatini*, *Hypernerphiini*, *Dociostaurini* etc.; endemics – *Ferganusa*, *Squamiana*, *Melanotmethis*, *Mizonocara* etc.). And so the distribution of the centres of diversity and endemism corresponds in outline to the general distribution of the *Orthoptera*.

The centres of generic diversity and species endemism are distributed similarly. But the degree of differentiation is higher. The significant barriers are observed in the steppe zone. This differentiation rises from north to south. There are many locally endemic species in the mountains, especially in Middle Asia. The main centres of endemism are connected with outlying ridges separated by the deep valleys: Gissar, Zaravshan, Turkistan, Kopetdag, Dzungarian Alatau. As a rule, the combination of the centres of diversity and the centers of endemism is generally observed. But in the plain or and the northern parts of North and Middle Asia the centres of diversity may be found independently of the corresponding centres of endemism (e.g. *Chorthippus*). On the contrary, we can observe specific centres of endemism unaccompanied by the corresponding diversity centres in isolated parts of the Earth's surface (e. g. islands).

**Acknowledgements.** I wish to express my sincere thanks to Prof. I. V. STEBAEV for his discussion of orthopteran biogeographical problems. I am also greatly indebted to Prof. MISTSHENKO, Dr. A. V. GOROCHOV and Dr. I. PODGORNAYA from the Zoological Institute (Sanct-Petersburg).

## REFERENCES

- EMELJANOV A. F. 1974. Proposals on the classification and nomenclature of areas. Entomol. Obozr., **53**: 497- 552.
- SERGEEV M. G. 1986. Patterns of *Orthoptera* distribution in North Asia. Nauka, Novosibirsk, 237 pp.
- SERGEEV M. G. 1988. Regularities of *Orthoptera* distribution in the mountains of the Middle Asia. Zool. Zh., **67**: 530-538.
- SERGEEV M. G. 1991. Distribution patterns of *Orthoptera* in the Asian part of the USSR. Thesis of doct. diss. Sanct-Petersburg. 37 pp.
- SERGEEV M. G. 1992. Distribution patterns of *Orthoptera* in North and Central Asia. J. Orth. Res., **1**: 14-24.
- SERGEEV M. G. 1993. Biogeography of Holarctic *Orthoptera*: faunas, populations, communities. Metaleptea, **14**(2): 11.
- UVAROV B. P. 1921. The geographical distribution of orthopterous insects in the Caucasus and in Western Asia. Proc. Zool. Soc. London, **31**: 447-472.